

# FATIGUE

**Carnitine** Transports fatty acids into mitochondria; Decreases both mental and physical fatigue in clinical trials.<sup>15,31,32</sup>

**Chromium** Promotes glucose uptake into cells, helping stabilize blood sugar.<sup>16,33</sup>

**Zinc** Deficiency lowers immunity and may cause muscle fatigue; Involved in several reactions for energy metabolism.<sup>15,34,35</sup>

**Asparagine** Supplementation of this amino acid delayed fatigue during exercise by decreasing the rate at which glycogen was used up; needed for gluconeogenesis, a process that allows glucose to be made from protein to prevent blood sugar from getting too low.<sup>1,2,3</sup>

**B Vitamins** Necessary for converting food into energy; Cofactors in the mitochondrial respiratory chain include B1, B2, B3, B5, B6, B12 and Folate.<sup>8,15,16,26-30</sup>

**Biotin** Helps liver utilize glycogen for energy. Animal studies confirm that biotin deficiency causes clinical fatigue.<sup>4</sup>

**Vitamin D** Low levels are seen in patients with chronic fatigue syndrome; Deficiency causes reduced muscle strength.<sup>24,25</sup>

**Glutamine** Mental and physical fatigue coincides with reduced levels of this amino acid in various tissues. Supplementation makes muscle more sensitive to insulin, increasing energy levels.<sup>5,6,7</sup>

**Vitamin E** Inverse correlation exists between fatigue and vitamin E levels.<sup>23</sup>

**Serine** Counteracts the overproduction of fatigue-causing stress hormones.<sup>8,9</sup>

**Vitamin A** When cellular levels of vitamin A are low, mitochondrial respiration and ATP production decreases.<sup>22</sup>

**CoQ10** Deficiency causes fatigue due to its role in mitochondrial energy metabolism; therapeutic benefits particularly noticeable in chronic fatigue syndrome.<sup>10,11,12,15</sup>

**Vitamin C** Assists iron uptake and transport; Precursor to carnitine and several hormones that affect energy levels. Supplementation reduced fatigue in various trials.<sup>15,16,21</sup>

**Antioxidants** Several studies confirm that oxidative stress exacerbates clinical symptoms of fatigue. Mitochondrial dysfunction (inefficient energy metabolism) can be treated therapeutically with antioxidants such as Selenium, Cysteine,  $\alpha$ -Lipoic acid and Glutathione, of which unusually low levels are seen in chronic fatigue patients.<sup>12,16,18,19,20</sup>

**Magnesium** Required to store energy molecule ATP; Repletion of magnesium in chronic fatigue patients shows clinical improvement in energy levels.<sup>15,16,17</sup>

**Fructose Intolerance** Fatigue (and hypoglycemia) are classic symptoms of this condition, since it depletes the main form of cellular energy, ATP.<sup>13,14</sup>

## REFERENCES

- <sup>1</sup>Sanpedro M, Roschel H et al. Effect of aspartate and asparagine supplementation on fatigue determinants in intense exercise. *Int J Sport Nutr Exerc Metab* 2013;13:65-75.
- <sup>2</sup>Lucas A, Rocco M et al. Effect of aspartate, asparagine, and carnitine supplementation in the diet on metabolism of skeletal muscle during a metabolic exercise. *Physiol Scholar* 1995;57:367-371.
- <sup>3</sup>Chivroux M, Phillips J, Berry M. Modulation of the inhibitory effect of chronic oxidation on gluconeogenesis from lactate by asparagine or less concentrations of aspartate. *Biochim Biophys Acta* 1995;1244:308-318.
- <sup>4</sup>Owens K, Kowal M et al. Experimental study of fatigue provoked by beta-alanine deficiency in mice. *Int J Vitam Nutr Res* 2014;74:334-340.
- <sup>5</sup>Jin G, Kikuchi Y et al. Changes in plasma and tissue amino acid levels in an animal model of complex fatigue. *Nutrition* 2009;25:520-527.
- <sup>6</sup>Prada P, Hoshino O et al. L-glutamine supplementation increases insulin resistance in adipose tissue and improves insulin signaling in liver and muscle of rats with diet-induced obesity. *Diabetologia* 2007;50:1949-1958.
- <sup>7</sup>Greenfield J, Frazee J et al. Oral glutamine increases circulating glucose-like peptide 1, glucagon, and insulin concentrations in lean, obese, and type 2 diabetic subjects. *Am J Clin Nutr* 2009;89:105-113.
- <sup>8</sup>Yoon K, Kelly G. Metabolic and behavioral correlates of chronic stress fatigue, neuroendocrine imbalance, anxiety, and restless sleep. *Aliment Pharmacol Ther* 2005;14:114-140.
- <sup>9</sup>Sturis M, Sturis S et al. The effects of phosphatidylserine on endocrine response to moderate intensity exercise. *J Int Soc Sports Nutr* 2008;5:11.
- <sup>10</sup>Mitsuru K, Tamaki M et al. Antifatigue effects of coenzyme Q10 during physical fatigue. *Nutrition* 2008;24:293-299.
- <sup>11</sup>Miles M, Mihalikova I et al. Coenzyme Q10 deficiency is synaptic encephalomyopathy / chronic fatigue syndrome (ME/CFS) is related to fatigue, autonomic and neurocognitive symptoms and to another risk factor explaining the early mortality in ME/CFS due to cancer. *Neuro Endocrinol Lett* 2009;30:47-476.
- <sup>12</sup>Logan A, Wang C. Chronic fatigue syndrome: oxidative stress and nitric oxide. *Aliment Pharmacol Ther* 2011;35:452-459.
- <sup>13</sup>Geary M, Douthett C et al. Doctor, my son is so tired... about a case of hereditary fructose intolerance. *Ann Endocrinol* 2007;18:455-458.
- <sup>14</sup>Liljestrand G, Hoptach E et al. Fructose transport and metabolism in adipose tissue of Zucker rats: diminished GLUT5 activity during obesity and insulin resistance. *Mol Cell Biochem* 2004;261:23-33.
- <sup>15</sup>Medbach M. Nutritional strategies for treating chronic fatigue syndrome. *Aliment Pharmacol Ther* 2000;5:53-108.
- <sup>16</sup>Hartmann E, Maggii G, Ruf M. The role of vitamins and minerals in energy metabolism and well-being. *J Int Med Res* 2007;35:277-289.
- <sup>17</sup>Moorhead G, Mansueti Y et al. Magnesium deficit in a sample of the Belgium population presenting with chronic fatigue. *Magnesium* 1997;18:329-337.
- <sup>18</sup>Miles M, Mihalikova I et al. Not in the mind of neuroendocrine hypotheses but in the cell nucleus: patients with chronic fatigue syndrome have increased production of nuclear factor kappa beta. *Neuro Endocrinol Lett* 2007;28:455-462.
- <sup>19</sup>Sturinus G, Molteni J. Competition for glutathione processes induces the immune system and the skeletal muscle: pathogenesis of chronic fatigue syndrome. *Mol Hypotheses* 1999;5:347-349.
- <sup>20</sup>Morales B, Cortés K. Reversing mitochondrial dysfunction, fatigue and the adverse effects of chemotherapy of metabolic disease by molecular replacement therapy. *Cell Exp Mol Biotechnol* 2008;25:161-168.
- <sup>21</sup>Qin S, Bao W et al. Inadequate vitamin C administration reduces fatigue in office workers: a double-blind randomized controlled trial. *Nutr J* 2012;11:7.
- <sup>22</sup>Pinto-Perez R, Hayes B et al. Effect of oxidative phosphorylation by vitamin A derivatives in mitochondrial rate in mitochondrial energy homeostasis. *FASEB J* 2010;24:557-566.
- <sup>23</sup>Vocchel J, Cluzette F et al. Relationship between neuroendocrine symptoms and blood markers of oxidative stress in patients with chronic fatigue syndrome. *Neurosci Lett* 2003;335:151-154.
- <sup>24</sup>Storchová B, Avstler B et al. Down-25-hydroxy Vitamin D Levels in Chronic Fatigue Syndrome: a Retrospective Study. *Int J Vitam Nutr Res* 2009;79:250-254.
- <sup>25</sup>Quirchuk L, Holtsch W. Vitamin D and reestablishing proper functional outcomes. *Nutr Clin Pract* 2007;22:287-304.
- <sup>26</sup>Hayes B, Polons T, Wessely S. Vitamin B status in patients with chronic fatigue syndrome. *J R Soc Med* 1999;92:183-185.
- <sup>27</sup>Quirchuk T, Hoshino H et al. Anti-fatigue effect of nicotinamide trypan blue is likely associated with excellent absorptivity and high bioavailability in tissues as a Vitamin B(1). *Eur J Pharmacol* 2010;635:117-123.
- <sup>28</sup>Vicenti S, Miron H et al. Thiamine (benfotiamine) stabilite improves energy metabolism and physical performance during physical-fatigue loading in rats. *Nutr Res* 2009;29:867-872.
- <sup>29</sup>Ergle A, Scherer U, Schreiner D. Efficacy of a combination therapy with vitamins B6, B12 and folic acid for general feeling of ill-health. Results of a non-interventional post-marketing surveillance study. *AGW Fortschkr Med* 2008;149 Suppl 4:162-165.
- <sup>30</sup>Laslett K, Quid S et al. Clinical activity of folic acid in patients with chronic fatigue syndrome. *Acute Medicine* 2006;5:399-404.
- <sup>31</sup>Mihalikova M, Gombart L et al. L-Carnitine treatment reduces severity of physical and mental fatigue and increases cognitive functions in cancer survivors: a randomized and controlled clinical trial. *Am J Clin Nutr* 2007;85:1738-1744.
- <sup>32</sup>Venkovski R, Schick H. Exploratory open label, randomized study of acetyl- and propionylcarnitine in chronic fatigue syndrome. *Psychosomatic Med* 2004;66:275-282.
- <sup>33</sup>Ortiz W, Reed J et al. Characterization of the metabolic and physiologic response to chromium supplementation in subjects with type 2 diabetes mellitus. *Diabetes* 2010;59:755-762.
- <sup>34</sup>Miles M, Mihalikova I et al. In chronic fatigue syndrome, the decreased levels of omega-3 polyunsaturated fatty acids are related to increased serum zinc and decrease in T cell activation. *Neuro Endocrinol Lett* 2005;26:745-751.
- <sup>35</sup>Castro A, Alvarez-Llan M. Behaviour of zinc in physical exercise: a special reference to immunity and fatigue. *Neurosci Biobehav Rev* 1995;19:439-445.

For more references, go to <http://www.AmericanJournalofNutrition.com/chronic-fatigue-syndrome/>